ATIENT NA	ME:	_ ID#: _		DATE:
escription: Tl	nis survey is meant to help us obtain information	on from our	patie	nts regarding their current levels of
scomfort and	capability. Please circle the answers below t	hat best ap	ply.	
. Please rate	e your pain level with activity: NO PAIN	=0 1 2	2 3	4 5 6 7 8 9 10 = VERY SEVERE PAIN
ODIFIED	OSWESTRY DISABILITY SCALE	– INITI	AL V	<u>/ISIT</u>
Pain Intensi			6.	Standing
) I can tolerate	the pain I have without having to use pain medicati	on.		I can stand as long as I want without increased pain.
) The pain is b pain medicat	ad, but I can manage without having to take		٠,,	, , , , ,
	ion provides me with complete relief from pain.		(2)	
) Pain medication	on provides me with moderate relief from pain.			Pain prevents me from standing more than 1/2 hour. Pain prevents me from standing more than 10 minutes.
	ion provides me with little relief from pain.		(5)	Pain prevents me from standing more than 10 minutes.
	ion has no effect on my pain.		(2)	Tam prevents me nom standing at an.
D	or (more delices described and a)		7.	Sleeping
	re (washing, dressing, etc.) re of myself normally without causing increased pair	n		Pain does not prevent me from sleeping well.
	e of myself normally, but it increases my pain.	1.		I can sleep well only by using pain medication. Even when I take pain medication, I sleep less than 6 hou
	o take care of myself, and I am slow and careful.			Even when I take pain medication, I sleep less than 4 hou
	out I am able to manage most of my personal care.		(4)	Even when I take pain medication, I sleep less than 2 hou
	very day in most aspects of my care.		(5)	Pain prevents me from sleeping at all.
) I do not get d	ressed, wash with difficulty, and stay in bed.			
Lifting			8.	Social Life
0	y weights without increased pain.		(0)	My social life is normal and does not increase my pain. My social life is normal, but it increases my level of pain.
	y weights, but it causes increased pain.			Pain prevents me from participating in more energetic
	me from lifting heavy weights off the floor,		(2)	activities (eg. sports, dancing).
	age if the weights are conveniently positioned		(3)	Pain prevents me from going out very often.
(eg, on a table			(4)	Pain has restricted my social life to my home.
	me from lifting heavy weights, but I can manage		(5)	I have hardly any social life because of my pain.
light to medii	um weights if they are conveniently positioned. very light weights.		0	Traveling
	or carry anything at all.		<b>9.</b> (0)	I can travel anywhere without increased pain.
, i cannot the c	r carry anything at an.			I can travel anywhere, but it increases my pain.
Walking				My pain restricts my travel over 2 hours.
	prevent me from walking any distance.			My pain restricts my travel over 1 hour.
	me from walking more than 1 mile.		(4)	My pain restricts my travel to short necessary journeys
	s me from walking more than ½ mile.		(5)	journeys under 1/2 hour.
Pain prevents	s me from walking more than 1/4 mile.  slk with crutches or a cane.		(5)	My pain prevents all travel except for visits to the physician/therapist or hospital.
	nost of the time and have to crawl to the toilet.			physician/therapist of nospital.
,			10.	Employment / Homemaking
Sitting			(0)	
	y chair as long as I like.		(1)	My normal homemaking/job activities increase my
	in my favorite chair as long as I like.		(2)	pain, but I can still perform all that is required of me.  1 can perform most of my homemaking/job duties, but
	s me from sitting more than 1 hour. s me from sitting more than ½ hour.		(2)	pain prevents me from performing more physically
	s me from sitting more than 10 minutes.			stressful activities (eg, lifting, vacuuming).
	s me from sitting at all.		(3)	Pain prevents me from doing anything but light duties.
			(4)	Pain prevents me from doing even light duties.
				Pain prevents me from performing any job or homemaking
			cho	res.
DIO lavarre	Fairbank 1080 All rights reserved ODI	ant info	rtion.	and narmission to use MADI December Tours
	contact@mapi-trust.org – Internet: <u>www.m</u>			and permission to use: MAPI Research Trust, Lyon
The second second				
Therapist Use C				
Comorbidities:		sorders (e.g., I	Parkins	on's, Muscular Dystrophy, Huntington's, CVA, Alzheimer's, TBI)
	☐ Diabetes ☐ Obesity ☐ Heart Condition ☐ Surgery for this	Problem		ICD9 Code:
			ıs, Rhei	umatoid Arthritis, Fibromyalgia)